

Malunion / Nonunion Management: What I Wish Someone Had Told Me Before I Started Doing These Cases

Preoperative assessment and types of nonunions

- I. Nonunion types
 - a. Hypertrophic
 - i. Characterized by good biology, poor stability
 - ii. “Horse’s hoof” appearance of fracture ends
 - iii. Bone formation obvious but persistent gap and/or motion at fracture site
 - iv. NEEDS STABILITY
 - b. Atrophic
 - i. Characterized by poor biology, stability unimportant
 - ii. Narrowed/sclerotic ends of fracture without obvious callus formation
 - iii. NEEDS BIOLOGY (and may need improved stability as well)
 - c. Oligotrophic
 - i. Part of a “spectrum of nonunion”...somewhere between atrophic and hypertrophic
 - ii. Likely retains biological capacity to heal but healing has not initiated
- II. Assessment of nonunions
 - a. Infection should always be suspected, especially after open fracture
 - i. History of problematic wound, recent drainage, prior infection
 - ii. C-reactive protein, erythrocyte sedimentation rate, white blood cell count (N.B. these may be normal even in an infected nonunion)
 - iii. (Consider indium-tagged WBC scan)
 - b. Patient-dependent factors
 - i. Comorbidities (diabetes, vascular disease)
 - ii. Tobacco or heavy ethanol use
 - iii. NSAIDs or other drugs that can slow bone healing (e.g. corticosteroids)
 - iv. Metabolic abnormalities (vitamin D deficiency, hypothyroidism, hypogonadism) – consider metabolic workup
 - v. Nutritional abnormalities (malnutrition due to behavior or disease)
 - vi. Genetics
 - c. Patient-independent factors
 - i. Injury characteristics (open fracture, bone loss, soft tissues)
 - ii. Surgery characteristics (distraction of fracture site, inadequate fixation)
 - iii. Infection (as above)

- d. Deformity
 - i. Plain radiographs are helpful for measuring maximal angular deformity of long bone nonunions/malunions
 - ii. Scanograms for suspected length deficiencies
 - iii. CT scan
 - 1. Fully delineate nonunions or malunions at the articular surface
 - 2. Bilateral extremity CT scans can assist with diagnosing rotational malunions or nonunions.
- e. Prior to any intervention, assess the overall picture
 - i. Would the surgical treatment be worse than the disease?
 - ii. Has sufficient time elapsed prior to diagnosing a nonunion?
 - 1. Tibial shaft fractures not healed at nine months...consider waiting longer (12 months)
 - iii. Have nonoperative modalities been exhausted?
 - 1. Optimize nutrition
 - 2. Bone stimulators (efficacy?)