

1. Calcaneous Fractures: Tips from the trenches
 - a. Patient
 - i. Modifiable factors
 1. Nicotine
 2. Diabetes
 - ii. Non Modifiable Factors
 1. Behavioral
 2. Work Status
 - b. Timing
 - i. What needs to be addresses acutely
 1. Skin at risk
 - a. Tongue type
 - b. Displaced Tuberosity: Joint Depression
 2. Open Injuries: Irrigate debride acutely. Wait for soft tissue envelope to mature before treating definitively
 - a. Medial
 - b. Plantar
 - c. Pattern
 - i. Tongue Type
 - ii. Joint Depression
 1. Medial column
 2. Lateral column
 3. Calcaneocuboid joint
 - d. How to address
 1. I&D
 2. Perc Pinning
 - a. Positioning
 - i. Lateral
 - ii. Prone
 - b. Approach
 - i. Perc
 - ii. Sinus Tarsi Incision
 - iii. Extensile Lateral (No- except with certain open injuries
 - iv. Clean out the sinus tarsi. You should be able to visualize the middle facet articulation with the talar neck.
 - c. Reduction
 - i. Schanz Pin
 - ii. Ex Fix
 - iii. Neuromuscular relaxation
 - iv. Threaded Steinman pin and curette. The tuberosity jack.
 - d. Implant
 - i. K-wire
 - ii. Cannulated Screws

- e. Stuff No One Ever Told Me
 - i. Wound management
 - 1. Closure
 - a. Make sure that the fascia is released from the tuberosity in confluence with the fascia overlying the short abductors. Close this with interrupted sutures which are placed snapped then tied. Close progressively toward the corner
 - b. Suture material
 - i. Braided absorbable pop-offs deep layer
 - ii. Monofilament absorbable (like Monocryl) superficial.
 - 1. Horizontal Mattress or Donatti
 - 2. Why? Because easy to leave in. No need to make the difficult decision when to remove.
 - ii. Acute Management
 - 1. Check the heel skin
 - a. Reduce the tuberosities urgently if skin at risk. Managing eschars is not what you want to do.
 - b. Elevate
 - c. Protect
 - d. Wait
 - iii. Get the anterior process right
 - a. Don't forget about the calcaneo-cuboid joint
 - b. It is fine to span the calcaneo-cuboid joint
 - iv. Consider bone substitutes
 - a. Biological Cement
 - b. Methylmethacrylate
 - c. PLGA c/allograft bone

Six tips learned the hard way.

1. Address compromised skin urgently
2. Reduce the calcaneocuboid joint. Span the joint if need be.
3. Visualize the middle facet articulation with the inferior talar neck. Confirms the overall quality of the reduction
4. Use biological cements or moldable graft to fill voids to reinforce your fixation.
5. Check the subtalar motion before you close. Again confirms quality of reduction and appropriate placement of hardware—nothing violating the subtalar joint.

6. Absorbable monofilament for final skin closure. Avoids wondering when to take the stitches out.

BIBLIOGRAPHY

[Treatment of displaced intra-articular calcaneal fractures with closed reduction and **percutaneous** screw fixation.](#)

Tomesen T, Biert J, Frölke JP.

J Bone Joint Surg Am. 2011 May 18;93(10):920-8.

[Minimally invasive treatment of calcaneal fracture by **percutaneous** leverage, anatomical plate, and compression bolts--the clinical evaluation of cohort of 156 patients.](#)

Wang Q, Chen W, Su Y, Pan J, Zhang Q, Peng A, Wu X, Wang P, Zhang Y.

J Trauma. 2010 Dec;69(6):1515-22.

[**Percutaneous** reduction and fixation of displaced intra-articular calcaneus fractures.](#)

DeWall M, Henderson CE, McKinley TO, Phelps T, Dolan L, Marsh JL.

J Orthop Trauma. 2010 Aug;24(8):466-72.

[A staged treatment plan for the management of Type II and Type IIIA open **calcaneus** fractures.](#)

Mehta S, Mirza AJ, Dunbar RP, Barei DP, Benirschke SK.

J Orthop Trauma. 2010 Mar;24(3):142-7.

[**Percutaneous** treatment of less severe intraarticular calcaneal fractures.](#)

Rammelt S, Amlang M, Barthel S, Gavlik JM, Zwipp H.

Clin Orthop Relat Res. 2010 Apr;468(4):983-90. Epub 2009 Jul 7.

[Complications when using threaded K-wire fixation for displaced intra-articular calcaneal fractures.](#)

Demcoe AR, Verhulsdonk M, Buckley RE.

Injury. 2009 Dec;40(12):1297-301. Epub 2009 Jun 16

[Limited open reduction and internal fixation of displaced intra-articular fractures of the calcaneum.](#)

Weber M, Lehmann O, Sägesser D, Krause F.
J Bone Joint Surg Br. 2008 Dec;90(12):1608-16.

[Secondary soft tissue compromise in tongue-type calcaneus fractures.](#)

Gardner MJ, Nork SE, Barei DP, Kramer PA, Sangeorzan BJ, Benirschke SK.

[Surgical treatment of intra-articular calcaneal fractures: a review of small incision approaches.](#)

Carr JB.
J Orthop Trauma. 2005 Feb;19(2):109-17. Review.